Author Index

Akira, K., see Baba, S. 251
Ames, M.M.
— , Miller, K.J. and Moertel, D.M.
High-performance liquid chromatographic assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 89
Amin, P.R., see Stetson, P.L. 217
Ang, K.P., see Gunasingham, H. 271
Aoki, I., see Tadano, K. 228
Arima, M., see Iwasaki, S. 182
Arita, T., see Tadano, K. 228
Baba, S.
— , Akira, K., Horie, M. and Mori, Y.
Application of radioisotope tracer techniques to analytical gas chromatography: determination of gas chromatographic peak yield 251
Baker, R.W.R., see Cullen, M.P. 420
Ballantyne, D.J., see Perrigo, B.J. 81
Barter, P.J., see Ha, Y.C. 154
Berthet, D., see Necciari, J. 202
Bosin, T.R.
— and Jarvis, C.A.
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by liquid chromatography 287
Bres, J., see Bressolle, F. 391
Bressolle, F.
— and Bres, J.
Dosage du sulpiride et du sultopride par chromatographie liquide à haute performance en vue de leur étude pharmacocinetique 391
Brown, A.S.
— , Cho, K.Y., Cheung, H.T.A., Hemmens, V. and Vine, J.
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary gas chromatography—mass spectrometry 139
Cautreels, W., see Necciari, J. 202
Cheung, H.T.A., see Brown, A.S. 139
Cho, K.Y., see Brown, A.S. 139
Chou, P.P.
— and Jaynes, P.K.
Determination of urinary 5-hydroxyindole-3-acetic acid using solid-phase extraction and reversed-phase high-performance liquid chromatography with electrochemical detection 167
Christ, D.D., see Walle, T. 213
Coombe, R.G.
— , Vine, J.H. and Yip, H.
Unusual fatty acids from amniotic fluid phospholipids 146
Crapper McLachlan, D.R., see Kruck, T.P.A. 123
Cuddy, K.K., see Skrabalak, D.S. 261
Cuisinaud, G.
— , Terrier, M., Ferry, N., Proust, S. and Sassard, J.
High-performance liquid chromatographic determination of cicletanide, a new diuretic, in plasma, red blood cells, urine and saliva 97
Cullen, M.P.
— and Baker, R.W.R.
Confidence limits for a gel-permeation system 420
Cummings, J.
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by high-performance liquid chromatography 401
Derendorf, H., see Yost, R.L. 131
De Silva, J.A.F., see Strojny, N. 313
Dixon, R., see Lucek, R. 239
Ensminger, W.D., see Stetson, P.L. 217
Ericsson, O., see Svensson, J.-O. 193
Esclade, L.
— , Guillochon, D. and Thomas, D.
Determination of metabolites of cytochrome P-450 model systems using high-performance liquid chromatography 373
Ferry, N., see Cuisinaud, G. 97
Foglietti, M.-J., see Succari, M. 457
Franklin, M., see Molyneux, S.G. 160
Fromark, I., see Johansson, B. 462
Fujiwara, S., see Suzuki, H. 341
Gombert, J., see Picard, M. 445
Guillochon, D., see Esclade, L. 373
Gunasingham, H.
— , Tay, B.T. and Ang, K.P.
Determination of oestriol in pregnancy urine by normal-phase high-performance liquid chromatography with electrochemical detection 271
Ha, Y.C. and Barter, P.J.
Rapid separation of plasma lipoproteins by gel permeation chromatography on agarose gel Superose 6B 154

Haginaka, J.
— , Wakai, J., Yasuda, H., Uno, T. and Nakagawa, T.
High-performance liquid chromatographic assay of sulbactam using pre-column reaction with 1,2,4-triazole 115
Hamada, C.
— , Iwasaki, M., Kuroda, N. and Ohkura, Y.
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in liquid chromatography and its use for the assay of 17-oxosteroids in urine 426
Hamilton, G.
— , Roth, E., Wallisch, E. and Tichy, F.
Semi-automated high-performance liquid chromatographic determination of cyclosporine A in whole blood using one-step sample purification and column-switching 411
Hansen, L.B., see Larsen, N.-E. 244
Hemmens, V., see Brown, A.S. 139
Hengy, H.
— and Kölle, E.-U.
Determination of gabapentin in plasma and urine by high-performance liquid chromatography and pre-column labelling for ultraviolet detection 473
Henion, J.D., see Skrabalak, D.S. 261
Henry, J.A., see Storey, G.C.A. 223
Hermens, W.A.J.J., see Overdiek, J.W.P.M. 279
Higuchi, S.
— , Urano, C. and Kawamura, S.
Determination of plasma protein binding of propafenone in rats, dogs and humans by highly sensitive gas chromatography—mass spectrometry 305
Hjertén, S.
— and Wu, B.-L.
Studies of fish zona pellucida by high-performance ion-exchange chromatography on agarose columns and free zone electrophoresis 295
Horie, M., see Baba, S. 251
Hsia, J.C., see Wong, L.T. 452
Hultin, T.A.
— , Mehta, R.G. and Moon, R.C.
Simple high-performance liquid chromatographic method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 187
Iwasaki, M., see Hamada, C. 426
Iwasaki, S.
— , Tanaka, H., Nakazawa, K. and Arima, M.
Quantitative high-performance liquid chromatography of bases and nucleosides in cerebral DNA of rat foetus 182
Jarvis, C.A., see Bosin, T.R. 287
Jaynes, P.K., see Chou, P.P. 167
Johansson, B.
— and Fromark, I.
Determination of meprobamate as an *n*-butylboronate ester derivative in serum by gas—liquid chromatography 462
Josephson, M.W., see Tsai, M.Y. 1
Kabra, P.M.
— and Nzekwe, E.U.
Liquid chromatographic analysis of clonazepam in human serum with solid-phase (Bond-Elut®) extraction 383
Kalow, W., see Kruck, T.P.A. 123
Kaneda, N.
— and Nagatsu, T.
Highly sensitive assay for choline acetyltransferase activity by high-performance liquid chromatography with electrochemical detection 23
Kawamura, S., see Higuchi, S. 305
Kempen, G.M.J. van, see Pennings, E.J.M. 172
Knodgen, B., see Seiler, N. 11
Knudsen, P., see Larsen, N.-E. 244
Kölle, E.U., see Hengy, H. 473
Koizumi, K.
— , Kubota, Y., Okada, Y. and Utamura, T.
Microanalyses of β-cyclodextrin in plasma by high-performance liquid chromatography 31
Kojima, K., see Suzuki, H. 176
Kondo, S., see Suzuki, H. 341
Kruck, T.P.A.
— , Kalow, W. and Crapper McLachlan, D.R.
Determination of desferoxamine and a major metabolite by high-performance liquid chromatography. Application to the treatment of aluminium-related disorders 123
Kubota, Y., see Koizumi, K. 31
Kuroda, N., see Hamada, C. 426
Kuroda, N., see Matsuoka, C. 432
Kurowski, M.
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair liquid chromatography and fluorometric detection 208
Lang, J.F., see Smithers, J.A. 232
Larsen, N.-E.
— , Hansen, L.B. and Knudsen, P.
Quantitative determination of perphenazine and its dealkylated metabolite

using high-performance liquid chromatography 244
Lempiäinen, M.
— and Mäkelä, A.-L.
Determination of proquazone and its *m*-hydroxy metabolite by high-performance liquid chromatography. Clinical application: pharmocokinetics of proquazone in children with juvenile rheumatoid arthritis 105
Lin, L.
— , Saller, C.F. and Salama, A.I.
Rapid automated high-performance liquid chromatographic analysis of cyclic adenosine 3′-5′-monophosphate. Synthesis in brain tissues 43
Lindberg, R.L.P.
Selective and sensitive high-performance liquid chromatographic assay for the metabolites of nomifensine in human plasma 333
Lintz, W.
— and Uragg, H.
Quantitative determination of tramadol in human serum by gas chromatography—mass spectrometry 65
Lucek, R.
— and Dixon, R.
Quantitation of levorphanol in plasma using high-performance liquid chromatography with electrochemical detection 239
McLachlan, D.R. Crapper, see Kruck, T.P.A. 123
Mäkelä, A.-L., see Lempiäinen, M. 105
Matsuoka, C.
— , Nohta, H., Kuroda, N. and Ohkura, Y.
Simultaneous determination of cholestanol and cholesterol in human serum by high-performance liquid chromatography with fluorescence detection 432
Mehta, R.G., see Hultin, T.A. 187
Merkus, F.W.H.M., see Overdiek, J.W.P.M. 279
Mery, D., see Necciari, J. 202
Miller, K.J., see Ames, M.M. 89
Miyazaki, K., see Tadano, K. 228
Moertel, D.M., see Ames, M.M. 89
Molyneux, S.G.
— and Franklin, M.
Routine determination of unconjugated 3-methoxy-4-hydroxyphenylglycol in plasma using high-performance liquid chromatography with electrochemical detection 160
Moon, R.C., see Hultin, T.A. 187
Moore, T., see Whelpton, R. 361
Mori, Y., see Baba, S. 251
Mould, G., see Read, J. 437
Muchtar, A., see Svensson, J.-O. 193
Nagatsu, T., see Kaneda, N. 23
Nagatsu, T., see Suzuki, H. 176
Nakagawa, T., see Haginaka, J. 115
Nakazawa, K., see Iwasaki, S. 182
Necciari, J.
— , Mery, D., Sales, Y., Berthet, D. and Cautreels, W.
Determination of propisomide, a new antiarrhythmic agent, in biological samples by gas chromatography with a thermionic detector 202
Nohta, H., see Matsuoka, C. 432
Norris, K.J., see Thompson, J.A. 349
Nzekwe, E.U., see Kabra, P.M. 383
Ohkura, Y., see Hamada, C. 426
Ohkura, Y., see Matsuoka, C. 432
Okada, Y., see Koizumi, K. 31
Okerholm, R.A., see Smithers, J.A. 232
Olichon, D., see Picard, M. 445
Oliphant, C., see Tsai, M.Y. 1
Overdiek, J.W.P.M.
— , Hermens, W.A.J.J. and Merkus, F.W.H.M.
Determination of the serum concentration of spironolactone and its metabolites by high-performance liquid chromatography 279
Parkinson, D.
Sensitive analysis of butyrophenone neuroleptics by high-performance liquid chromatography with ultraviolet detection at 254 nm 465
Peel, H.W., see Perrigo, B.J. 81
Pennings, E.J.M.
— , Verhagen, J.C.M. and Van Kempen, G.M.J.
Assay of urinary phenylacetic acid by high-performance liquid chromatography 172
Percheron, F., see Succari, M. 457
Perrigo, B.J.
— , Peel, H.W. and Ballantyne, D.J.
Use of dual-column fused-silica capillary gas chromatography in combination with detector response factors for analytical toxicology 81
Petersen, D.R., see Thompson, J.A. 349
Picard, M.
— , Olichon, D. and Gombert, J.
Determination of serotonin in plasma by liquid chromatography with electrochemical detection 445
Proust, S., see Cuisinaud, G. 97

Read, J.
— , Mould, G. and Stevenson, D.
Simple high-performance liquid chromatographic method for the determination of medroxyprogesterone acetate in human plasma 437
Rhys Williams, A.T.
Simultaneous determination of serum vitamin A and E by liquid chromatography with fluorescence detection 198
Roth, E., see Hamilton, G. 411
Salama, A.I., see Lin, L. 43
Sales, Y., see Necciari, J. 202
Saller, C.F., see Lin, L. 43
Sassard, J., see Cuisinaud, G. 97
Schootstra, R., see Storey, G.C.A. 223
Seiler, N.
— and Knodgen, B.
Determination of amino acids by separation of their ion pairs with dodecyl sulphate 11
Shintani, H.
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53
Shukla, U.A., see Stetson, P.L. 217
Silva, J.A.F. de, see Strojny, N. 313
Skrabalak, D.S.
— , Cuddy, K.K. and Henion, J.D.
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-liquid chromatography—mass spectrometry 261
Smithers, J.A.
— , Lang, J.F. and Okerholm, R.A.
Quantitative analysis of vigabatrin in plasma and urine by reversed-phase high-performance liquid chromatography 232
Stetson, P.L.
— , Shukla, U.A., Amin, P.R. and Ensminger, W.D.
High-performance liquid chromatographic method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 217
Stevenson, D., see Read, J. 437
Storey, G.C.A.
— , Schootstra, R. and Henry, J.A.
Measurement of meptazinol in plasma by high-performance liquid chromatography with electrochemical detection 223
Strojny, N.
— and De Silva, J.A.F.
Determination of diclofensine, and antidepressant agent, and its major metabolites in human plasma by high-performance liquid chromatography with fluorometric detection 313
Succari, M.
— , Foglietti, M.-J. and Percheron, F.
Two-step purification of human α_1-acid glycoprotein 457
Sugimoto, I., see Suzuki, H. 341
Suzuki, H.
— , Fujiwara, S., Kondo, S. and Sugimoto, I.
Determination of nifedipine in human plasma by high-performance liquid chromatography with electrochemical detection 341
Suzuki, H.
— , Yata, J., Kojima, K. and Nagatsu, T.
Simple and sensitive assay of dopamine β-hydroxylase in human cerebrospinal fluid by high-performance liquid chromatography with electrochemical detection 176
Svensson, J.-O.
— , Muchtar, A. and Ericsson, Ö.
Ion-pair high-performance liquid chromatographic determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 193
Tadano, K.
— , Yuhki, Y., Oki, I., Miyazaki, K. and Arita, T.
High-performance liquid chromatographic determination of tranilast in plasma 228
Tanaka, H., see Iwasaki, S. 182
Tay, B.T., see Gunasingham, H. 271
Terrier, M., see Cuisinaud, G. 97
Thomas, D., see Esclade, L. 373
Thompson, J.A.
— , Norris, K.J. and Petersen, D.R.
Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1′-N-oxide formation in mice 349
Tichy, F., see Hamilton, G. 411
Tsai, M.Y.
— , Oliphant, C. and Josephson, M.W.
Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary gas chromatography 1

Uno, T., see Haginaka, J. 115
Uragg, H., see Lintz, W. 65
Urano, C., see Higuchi, S. 305
Utamura, T., see Koizumi, K. 31
Van Kempen, G.M.J., see Pennings, E.J.M. 172
Verhagen, J.C.M., see Pennings, E.J.M. 172
Vine, J., see Brown, A.S. 139
Vine, J.H., see Coombe, R.G. 146
Wakai, J., see Haginaka, J. 115
Walle, T.
— , Christ, D.D., Walle, U.K. and Wilson, M.J.
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by high-performance liquid chromatography after chiral derivatization 213
Walle, U.K., see Walle, T. 213
Wallisch, E., see Hamilton, G. 411
Whelpton, R.
— and Moore, T.
Sensitive liquid chromatographic method for physostigmine in biological fluids using dual-electrode electrochemical detection 361
Williams, A.T. Rhys, see Rhys Williams, A.T. 198
Wilson, M.J., see Walle, T. 213
Wong, L.T.
— , Xu, X.J. and Hsia, J.C.
Fractionation of rat α-fetoprotein by high-performance liquid chromatography 452
Wu, B.-L., see Hjertén, S. 295
Xu, X.J., see Wong, L.T. 452
Yasuda, H., see Haginaka, J. 115
Yata, J., see Suzuki, H. 176
Yip, H., see Coombe, R.G. 146
Yost, R.L.
— and Derendorf, H.
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 131
Yuhki, Y., see Tadano, K. 228

Subject Index

Acetylcholine
Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 23

Acetylisoniazid
Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 193

α_1-Acid glycoprotein
Two-step purification of human α_1-acid glycoprotein 457

Actinomadura R39
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC—MS 139

Adenosine 3',5'-monophosphate, cyclic
Rapid automated HPLC analysis of cyclic adenosine 3',5'-monophosphate. Synthesis in brain tissues 43

Amines, tertiary
Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1'-N-oxide formation in mice 349

Amino acids
Determination of amino acids by separation of their ion pairs with dodecyl sulphate 11

Bases
Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 182

Betamethasone
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC—MS 261

Bisbenzimidazole
HPLC assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 89

Bromodeoxyuridine
HPLC method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 217

Bromouracil
HPLC method for the determination of bromodeoxyuridine and its major metabolite, bromouracil, in biological fluids 217

Butyrophenones
Sensitive analysis of butyrophenone neuroleptics by HPLC with ultraviolet detection at 254 nm 465

Calcium
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53

Canrenone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 279

Cations
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53

Cefotaxime
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 131

3-Chloroformyl-7-methoxycoumarin
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 426

1-[γ-(2-Chloro-10-phenothiazinyl)propyl]-piperazine dihydrochloride
Quantitative determination of perphenazine and its dealkylated metabolite using HPLC 244

Cholestanol
Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 432

Cholesterol
Simultaneous determination of cholestanol and cholesterol in human serum by HPLC with fluorescence detection 432

Choline
Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 23

Choline acetyltransferase
Highly sensitive assay for choline acetyltransferase activity by HPLC with electrochemical detection 23

Cicletanide
HPLC determination of cicletanide, a new diuretic, in plasma, red blood cells, urine and saliva 97

Clonazepam
LC analysis of clonazepam in human serum with solid-phase (Bond-Elut®) extraction 383

β-Cyclodextrin
Microanalyses of β-cyclodextrin in plasma by HPLC 31
Cyclosporine A
Semi-automated HPLC determination of cyclosporine A in whole blood using one-step sample purification and column-switching 411
Cytochrome P-450
Determination of metabolites of cytochrome P-450 model systems using HPLC 373
4′-Deoxydoxorubicin
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
4′-Deoxydoxorubicin aglycone
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
4′-Deoxydoxorubicin 7-deoxyaglycone
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
4′-Deoxydoxorubicinol
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
4′-Deoxydoxorubicinol aglycone
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
4′-Deoxydoxorubicinol 7-deoxyaglycone
Method for the determination of 4′-deoxydoxorubicin, 4′-deoxydoxorubicinol and their 7-deoxyaglycones in human serum by HPLC 401
Desacetylcefotaxime
Rapid chromatographic determination of cefotaxime and its metabolite in biological fluids 131
Desferoxamine
Determination of desferoxamine and a major metabolite by HPLC. Application to the treatment of aluminium-related disorders 123
Desferoxamine aluminum(III) chelate
Determination of desferoxamine and a major metabolite by HPLC. Application to the treatment of aluminium-related disorders 123
Desferoxamine iron(III) chelate
Determination of desferoxamine and a major metabolite by HPLC. Application to the treatment of aluminium-related disorders 123
Detector response factors
Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 81
Dextrans
Confidence limits for a gel-permeation system 420
Dextrose
Confidence limits for a gel-permeation system 420
Diclofensine
Determination of diclofensine, an antidepressant agent, and its major metabolites in human plasma by HPLC with fluorometric detection 313
DNA
Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 182
Dodecyl sulphate
Determination of amino acids by separation of their ion pairs with dodecyl sulphate 11
Dopamine β-hydroxylase
Simple and sensitive assay of dopamine β-hydroxylase in human cerebrospinal fluid by HPLC with electrochemical detection 176
Fatty acids
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC—MS 139
Fatty acids
Unusual fatty acids from amniotic fluid phospholipids 146
α-Fetoprotein
Fractionation of rat α-fetoprotein by HPLC 452
Gabapentin
Determination of gabapentin in plasma and urine by HPLC and pre-column labelling for ultraviolet detection 473
Haloperidol
Sensitive analysis of butyrophenone neuroleptics by HPLC with ultraviolet detection at 254 nm 465
Hexadecane
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 251

Histidine
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 251
6β-Hydroxybetamethasone
Quantitative determination of betamethasone and its major metabolite in equine urine by micro-LC—MS 261
5-Hydroxyindole-3-acetic acid
Determination of urinary 5-hydroxyindole-3-acetic acid using solid-phase extraction and reversed-phase HPLC with electrochemical detection 167
3′-Hydroxy-4′-methoxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 333
4′-Hydroxy-3′-methoxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 333
4′-Hydroxynomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 333
N-(4-Hydroxyphenyl)-all-*trans*-retinamide
Simple HPLC method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 187
4′-Hydroxypropranolol
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 213
4′-Hydroxypropranolol sulphate
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 213
6β-Hydroxy-7α-thiomethylspirolactone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 279
Indole ethylamines
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 287
Isoniazid
Ion-pair HPLC determination of isoniazid and acetylisoniazid in plasma and urine. Application for acetylator phenotyping 193
Ketanserin
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair LC and fluorometric detection 208
Ketanserinol
Simultaneous determination of ketanserin and ketanserinol in biological fluids using ion-pair LC and fluorometric detection 208
Levorphanol
Quantitation of levorphanol in plasma using HPLC with electrochemical detection 239
Lipoproteins
Rapid separation of plasma lipoproteins by gel permeation chromatography on agarose gel Superose 6B 154
Magnesium
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53
Medroxyprogesterone acetate
Simple HPLC method for the determination of medroxyprogesterone acetate in human plasma 437
Meprobamate
Determination of meprobamate as an *n*-butylboronate ester derivative in serum by GLC 462
Meptazinol
Measurement of meptazinol in plasma by HPLC with electrochemical detection 223
3-Methoxy-4-hydroxyphenylglycol
Routine determination of unconjugated 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 160
2-Methyl-2-propyl-1,3-propanediol *n*-butylboronate
Determination of meprobamate as an *n*-butylboronate ester derivative in serum by GLC 462
Nicotine-1′-N-oxide
Isolation and analysis of N-oxide metabolites of tertiary amines: quantitation of nicotine-1′-N-oxide formation in mice 349
Nifedipine
Determination of nifedipine in human plasma by HPLC with electrochemical detection 341
Nomifensine
Selective and sensitive HPLC assay for the metabolites of nomifensine in human plasma 333

Nordiclofensine
Determination of diclofensine, an antidepressant agent, and its major metabolites in human plasma by HPLC with fluorometric detection 313
Nucleosides
Quantitative HPLC of bases and nucleosides in cerebral DNA of rat foetus 182
Oestriol
Determination of oestriol in pregnancy urine by normal-phase HPLC with electrochemical detection 271
Organic acids
Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary GC 1
Organic acidurias
Identification of metabolites diagnostic for organic acidurias by simultaneous dual-column capillary GC 1
17-Oxosteroids
3-Chloroformyl-7-methoxycoumarin as a fluorescent derivatization reagent for alcoholic compounds in LC and its use for the assay of 17-oxosteroids in urine 426
Peak yields
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 251
Perphenazine
Quantitative determination of perphenazine and its dealkylated metabolite using HPLC 244
Phenylacetic acid
Assay of urinary phenylacetic acid by HPLC 172
Phospholipids
Unusual fatty acids from amniotic fluid phospholipids 146
Physostigmine
Sensitive LC method for physostigmine in biological fluids using dual-electrode electrochemical detection 361
Pibenzimol
HPLC assay and preclinical pharmacological studies of pibenzimol (bisbenzimidazole) 89
Potassium
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53
Prenalterol sulphate
Separation of the enantiomers of intact sulfate conjugates of adrenergic drugs by HPLC after chiral derivatization 213
Propafenone
Determination of plasma protein binding of propafenone in rats, dogs and humans by highly sensitive GC—MS 305
Propisomide
Determination of propisomide, a new antiarrhythmic agent, in biological samples by GC with a thermionic detector 202
Proquazone
Determination of proquazone and its *m*-hydroxy metabolite by HPLC. Clinical application: pharmacokinetics of proquazone in children with juvenile rheumatoid arthritis 105
Retention indices
Use of dual-column fused-silica capillary GC in combination with detector response factors for analytical toxicology 81
Retinoids
Simple HPLC method for the separation of retinoids including N-(4-hydroxyphenyl)-all-*trans*-retinamide 187
Serotonin
Determination of serotonin in plasma by LC with electrochemical detection 445
Sodium
Improvement of ion chromatography with ultraviolet photometric detection and comparison with conductivity detection for the determination of serum cations 53
Spironolactone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 279
Spiroperidol
Sensitive analysis of butyrophenone neuroleptics by HPLC with ultraviolet detection at 254 nm 465
Streptomyces R61
Determination of fatty acids of the bacteria *Streptomyces* R61 and *Actinomadura* R39 by capillary GC—MS 139
Sulbactam
HPLC assay of sulbactam using precolumn reaction with 1,2,4-triazole 115

Sulpiride
Quantitative analysis of sulpiride and sultropide by HPLC for pharmacokinetic studies 391
Sultopride
Quantitative analysis of sulpiride and sultopride by HPLC for pharmacokinetic studies 391
Testosterone
Application of radioisotope tracer techniques to analytical GC: determination of GC peak yield 251
Tetrahydro-β-carbolines
Derivatization in aqueous solution, isolation and separation of tetrahydro-β-carbolines and their precursors by LC 287
7α-Thiomethylspirolactone
Determination of the serum concentration of spironolactone and its metabolites by HPLC 279
Tramadol
Quantitative determination of tramadol in human serum by GC—MS 65
Tranilast
HPLC determination of tranilast in plasma 228
1,2,4-Triazole
HPLC assay of sulbactam using pre-column reaction with 1,2,4-triazole 115
Trifluperidol
Sensitive analysis of butyrophenone neuroleptics by HPLC with ultraviolet detection at 254 nm 465
Vigabatrin
Quantitative analysis of vigabatrin in plasma and urine by reversed-phase HPLC 232
Vitamin A
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 198
Vitamin E
Simultaneous determination of serum vitamin A and E by LC with fluorescence detection 198
Zona pellucida
Studies of fish zona pellucida by high-performance ion-exchange chromatography on agarose columns and free zone electrophoresis 295